Contents

Articles

1 Budd, G.E.: The morphology of *Opabinia regalis* and the reconstruction of the arthropod stem-group
15 Ramsköld, L., Chen, J.-y., Edgecombe, G.D. & Zhou, G.-q.: Preservational folds simulating tergite junctions in tegopeltid and naraoiid arthropods
21 Dattilo, B.F.: A quantitative paleoecological approach to high-resolution cyclic and event stratigraphy: the Upper Ordovician Miamitown Shale in the type Cincinnatian
39 Melchin, M.J. & Doucet, K.M.: Modelling flow patterns in conical dendroid graptolites
47 Baumiller, T.K. & Ausich, W.I.: Crinoid stalk flexibility: theoretical predictions and fossil stalk postures
61 Checa, A.: Origin of intracameral sheets in ammonoids
77 Boomer, I., Whatley, R. & Aladin, N.V.: Aral Sea Ostracoda as environmental indicators
87 Casinos, A.: Bipedalism and quadrupedalism in *Megatherium*: an attempt at biomechanical reconstruction
97 Werdelin, L.: Community-wide character displacement in Miocene hyaenas
113 Kowalewski, M. & Demko, T.M.: Trace fossils and population paleoecology: comparative analysis of size-frequency distributions derived from burrows
125 Heikoop, J.M., Tsujita, C.J., Heikoop, C.E., Risk, M.J. & Dickin, A.P.: Effects of volcanic ashfall recorded in ancient marine communities: comparison of a nearshore and an offshore environment
141 Savarese, M., Dodd, J.R. & Lane, N.G.: Taphonomic and sedimentologic implications of crinoid intraskeletal porosity
157 Mapes, R.H. & Mapes, G.: Biotic destruction of terrestrial plant debris in the Late Paleozoic marine environment
171 Watkins, R.: Paleoecology of Silurian reef bivalves, Racine Formation, North America
181 Aberhan, M. & Fürsich, F.T.: Diversity analysis of Lower Jurassic bivalves of the Andean Basin and the Pliensbachian–Toarcian mass extinction
197 Risk, M.J., Sayer, B.G., Tevesz, M.J.S. & Karr, C.D.: Comparison of the organic matrix of fossil and recent bivalve shells
203 Marshall, J.D., Pirrie, D., Clarke, A., Nolan, C.P & Sharman, J.: Stable-isotopic composition of skeletal carbonates from living Antarctic marine invertebrates
213 Kruse, P.D.: Hyolith guts in the Cambrian of northern Australia – turning hyolithomorphs upside down
219 Lethiers, F., Damotte, R. & Whatley, R.: Evidence of brooding in Permian non-marine Ostracoda
225 Checa, A., Company, M., Sandoval, J. & Weitschat, W.: Covariation of morphological characters in the Triassic ammonoid Czekanowskites rieberi
237 Ash, S.: Evidence of arthropod–plant interactions in the Upper Triassic of the southwestern United States
249 Kielan-Jaworowska, Z.: Characters of multituberculates neglected in phylogenetic analyses of early mammals
267 Doyle, P., Mather, A.E., Bennett, M.R & Bussell, M.A.: Miocene barnacle assemblages from southern Spain and their palaeoenvironmental significance
275 Edinger, E.N. & Risk, M.J.: Sponge borehole size as a relative measure of bioerosion and paleoproductivity
289 Miller, K.B. & West R.R.: Growth-interruption surfaces within chaetetid skeletons: Records of physical disturbance and depositional dynamics
301 Lenz, A.C. & Melchin, M.J.: Phylogenetic analysis of the Silurian Retiolitidae
311 Jeppsson, L.: Recognition of a probable secundo–primo event in the Early Silurian
317 Pridmore, P.A., Barwick, R.E. & Nicoll, R.S.: Soft anatomy and the affinities of conodonts
329 Girard, C. & Feist, R.: Eustatic trends in conodont diversity across the Frasnian–Famennian boundary in the stratotype area, Montagne Noire, Southern France
339 Mutvei, H.: Characterization of actinoceratoid cephalopods by their siphuncular structure
349 Williams, A. & Cusack, M.: Lingulid shell mediation in clay formation
361 Wendt, J.: Aragonitic dasycladalean algae from the Upper Permian of Sichuan, China
369 Krassilov, V.A. & Rasnitsyn, A.P.: Pollen in the guts of Permian insects: first evidence of pollinivory and its evolutionary significance
373 Unwin, D.M.: Pterosaur tracks and the terrestrial ability of pterosaurs

Lethaia Seminar

85 Peña, A. de la & Soler-Gijón, R.: The first siluriform fish from the Cretaceous–Tertiary interval of Eurasia
107 Towe, K.M.: Fossil preservation in the Burgess Shale
109 Butterfield, N.J.: Fossil preservation in the Burgess Shale: Reply
287 Torres, A.M.: Fossil algae were very different from xenophyophores
299 Evans, K.R. & Maybury, C.A.: Pennsylvanian phylloid algae interpreted as shallow-water xenophyophores: Reply

Lethaia Forum

38 Franzén, C. & Bengtson, S.: Heinz Lowenstam's collection of fossils
218 Lane, H.R. & Bruton, D.L.: *International Palaeontological Association*: Endangered fossil sites
236 Foster, M.W.: Fossil fish from Mazon Creek Biota stolen
310 Bertling, M.: 'Endolithos' – First International Workshop on Bioerosion, 2nd–7th September, 1996

Lethaia Reviews

139 Campbell, K.S.W.: Early vertebrates
170 Ahlberg, P.E.: *Ichthyostega* in depth
224 Weishampel, D.B.: Herbivory and reptiles